

## Steatoda Spiders

Family: Theridiidae

The Steatoda belongs to the family Theridiidae, also called the comb-footed spiders (Fig. 1). Steatoda is a common house spider that lives up to six years and is not known to be particularly harmful to people. Another important species of this family is the black widow spider, Latrodectus mactans. These spiders have two distinctive differences. Steatoda does not have the reddish orange hourglass shape on the ventral side of the abdomen and the shape of the abdomen is more oval and not as globose as in Latrodectus. It does however have an outline that resembles an hourglass on the ventral side of the abdomen. Steatoda is further distinguished from the widow in that it bears a light-colored crescent or band on the leading edge of the dorsal abdominal surface. Some Steatoda have been reported to actually prey on black widow spiders and cockroaches. A major food source is sowbugs.

## **Selected References:**

Borror, Donald J. and Dwight M. DeLong. 1971. *An Introduction to the Study of Insects*, 3<sup>rd</sup> ed. Holt, Rinehart, and Winston, Inc., NY, 811 p.

Ebeling, Walter. 1978. *Urban Entomology.* Division of Agricultural Sciences, University of California Berkeley. 694 p.

Kasten, J.B. 1972. *How to Know the Spiders*, 3<sup>rd</sup> ed. Wm. C. Brown Co. Publishers, Dubuque, Iowa. 272 p.



Fig. 1. A theridiid spider (*Steatoda* sp.)

By Brett M. Johnson